

---

# ULTRASTRUCTURAL STUDY OF CONIDIA OF *Colletotrichum gloeosporioides* AND *Colletotrichum musae* TREATED WITH ESSENTIAL OILS

---

Luciane Cristina Rozwalka, Eduardo Alves and Douglas Carvalho do Amaral

## SUMMARY

Essential oils have shown to be efficient in the control of plant diseases; however, no reports exist regarding their mode of action on plant pathogens. The aim of this work was to evaluate the effect of essential oils from *Cymbopogon martinii*, *Eugenia caryophyllata*, *Thymus vulgaris*, *Cinnamomum sp.* and *Cymbopogon citratus* on conidia of *Colletotrichum gloeosporioides* and *Colletotrichum musae*, etiologic agents of anthracnose of guava and banana, respectively, by means of transmission electron microscopy (TEM). Conidia suspensions ( $1 \times 10^{10}$  conidia/ml) prepared in sterile distilled water with Tween 20® 1.0% were treated with essential oils at 0.5%, remaining under agitation at 25°C for 24h. Water alone was

used as control. After centrifugation the supernatant was discarded and the masses of conidia obtained were fixed for 24h in modified Karnovsky fixative. The suspensions were centrifuged again and after discarding the supernatant, the fixed conidia were embedded in agarose gel and subjected to the protocol of sample preparation for TEM, to be observed with a Zeiss EM 109 microscope. The essential oils showed fungitoxic action directly on the conidia of *C. gloeosporioides* and *C. musae*, causing severe damage by promoting cellular disorganization and degradation that makes germination unviable.

---

## Introduction

The anthracnose caused in avocado, guava, papaya, mango and passion fruit by *Colletotrichum gloeosporioides* and in banana by *Colletotrichum musae* constitutes a major postharvest problem (Peres *et al.*, 2002). The essential oils have shown to be efficient in controlling *C. gloeosporioides* (Lee *et al.*, 2007; Barreira-Necha *et al.*, 2008; Duamkhanmanee, 2008; Rozwalka *et al.*, 2008; Sukatta *et al.*, 2008; Anaruma *et al.*, 2010) and other species of the genus (Ranasinghe *et al.*, 2002; Singh *et al.*, 2002; Shahi *et al.*, 2003; Arroyo *et al.*, 2007; Tzortzakakis and Economakis, 2007; Tzortzakakis, 2009). However, there are no reports regarding the mode of action of these on

plant pathogens (Arroyo *et al.*, 2007).

Due to the complexity of essential oils, it is assumed that there are multiple mechanisms of action, not well known, that could result on pathogen inhibition, such as protein denaturation, enzyme inhibition and/or membrane disintegration (Janssen, 1989 *apud* Siani *et al.*, 2000).

This study aimed to evaluate the mode of action of the essential oils of *Cymbopogon martinii*, *Eugenia caryophyllata*, *Thymus vulgaris*, *Cinnamomum sp.* and *Cymbopogon citratus* on conidia of *Colletotrichum gloeosporioides* and *Colletotrichum musae*, etiologic agents of the anthracnose in guava and banana, respectively, by means of transmission electron microscopy (TEM).

## Materials and Methods

### Place

The experiments were performed at the Laboratory of Electron Microscopy and Ultrastructural Analysis, Department of Plant Pathology, Universidade Federal de Lavras, Minas Gerais State, Brazil.

### Acquisition and maintenance of *Colletotrichum gloeosporioides* and *C. musae* isolates

The strains of *C. gloeosporioides* and *C. musae* were obtained from lesions on ripe fruits of guava (cv. Pedro Sato) and banana (cv. Prata), respectively, purchased in the local market and from local producers. After verifying the pathoge-

nicity, the isolates were maintained in potato dextrose agar (PDA), in growth chambers at 25°C and a 12h photoperiod.

### Treatments and sample preparation for TEM

For the evaluation of the mode of action of the essential oils on the ultrastructure of *C. gloeosporioides* and *C. musae*, conidia were subjected to treatments containing the essential oils of *Cinnamomum sp.* (cinnamon), *Cymbopogon citratus* (lemongrass), *Eugenia caryophyllata* (Indian clove), *Cymbopogon martinii* (palmarosa), and *Thymus vulgaris* (thyme), selected in function of the potential for total inhibition (100%) on the germination of the pathogens at concentrations of 0.1 and

---

## KEYWORDS / Alternative Control / Anthracnose / Antifungal Activity / Essential Oils / Transmission Electron Microscopy /

Received: 07/13/2010. Modified: 10/03/2010. Accepted: 10/15/2010.

Luciane Cristina Rozwalka. Agricultural Engineer and M.Sc. in Agronomy - Vegetal Production, Federal University of Paraná, Brazil. Doctor in Agronomy - Phytopathology, Federal University of Lavras (UFLA), Brazil. e-mail: luciane.rozwalka@gmail.com

Eduardo Alves. Agricultural Engineer and M.Sc. in Agronomy - Phytopathology, UFLA, Brazil. Doctor in Agronomy - Phytopathology, Superior School of Agriculture "Luiz de Queiroz", University of São Paulo, Brazil. Professor, UFLA, Brazil. Address: De-

partment of Plant Pathology, UFLA. 37200-000 - Lavras, MG, Brazil. P.O.Box: 3037. e-mail: ealves@dfp.ufla.br  
Douglas Carvalho do Amaral. Biology student, Center University of Lavras (UNILAVRAS), Brazil. Fellow, Research Foundation of the State of Minas

Gerais (FAPEMIG), Laboratory of Electron Microscopy and Ultrastructural Analysis, UFLA, Brazil. e-mail: douglaskamara@gmail.com

## ESTUDIO ULTRAESTRUCTURAL DE LAS CONIDIAS DE *Colletotrichum gloeosporioides* Y *Colletotrichum musae* TRATADAS CON ACEITES ESENCIALES

Luciane Cristina Rozwalka, Eduardo Alves y Douglas Carvalho do Amaral

### RESUMEN

Los aceites esenciales han demostrado eficacia en el control de enfermedades de las plantas; sin embargo, no existen reportes del modo de acción de estos sobre los fitopatógenos. El objetivo de este trabajo fue evaluar el efecto de los aceites esenciales de *Cymbopogon martinii*, *Eugenia caryophyllata*, *Thymus vulgaris*, *Cinnamomum sp.* y *Cymbopogon citratus* sobre los conidios de *Colletotrichum gloeosporioides* y *Colletotrichum musae*, agentes etiológicos de la antracnosis de la guayaba y del plátano, respectivamente, por medio de microscopía electrónica de transmisión (MET). Una suspensión de conidios ( $1 \times 10^{10}$  conidios/ml) preparada con agua destilada estéril y con Tween 20® 1,0%, fue tratada con aceites esenciales al 0,5%, mante-

niéndose en agitación a 25°C durante 24h. El control consistió de agua solamente. Después de la centrifugación y el descarte del sobrenadante, las masas de conidios obtenidas fueron fijadas por 24h en fijador de Karnovsky modificado. Las suspensiones se centrifugaron de nuevo y después de descartar el sobrenadante, las conidios fijadas fueron incluidas en gel de agarosa y sometidas al protocolo de preparación de muestras para MET. Utilizando un microscopio Zeiss EM 109 se observó que los aceites esenciales ejercieron una acción fungitóxica directa sobre las conidios de *C. gloeosporioides* y *C. musae*, causándoles daños notorios, promoviendo la desorganización y degradación celular que imposibilita la germinación.

## ESTUDO ULTRAESTRUTURAL DE CONÍDIOS DE *Colletotrichum gloeosporioides* E *Colletotrichum musae* TRATADOS COM ÓLEOS ESSENCIAIS

Luciane Cristina Rozwalka, Eduardo Alves e Douglas Carvalho do Amaral

### RESUMO

Os óleos essenciais têm demonstrado eficiência no controle de doenças de plantas; entretanto, praticamente inexistem relatos do modo de ação desses sobre fitopatógenos. O objetivo do trabalho foi avaliar o efeito de óleos essenciais de *Cymbopogon martinii*, *Eugenia caryophyllata*, *Thymus vulgaris*, *Cinnamomum sp.* e *Cymbopogon citratus* sobre conidios de *Colletotrichum gloeosporioides* e *Colletotrichum musae*, agentes etiológicos da antracnose em goiabas e bananas, respectivamente, por meio de microscopia eletrônica de transmissão (MET). Conídios, em suspensão ( $1 \times 10^{10}$  conídios/ml) preparada em água destilada e esterilizada com Tween 20® a 1,0%, foram tratados com os óleos essenciais a 0,5% permanecendo sob

agitação a 25°C, por 24 horas. No controle utilizou-se apenas água. Após centrifugação e descarte do sobrenadante, as massas de conídios obtidas foram fixadas por 24h em fixador de Karnovsky modificado. As suspensões foram novamente centrifugadas e após o descarte do sobrenadante, os conídios fixados foram emblocados em gel de agarose e submetidos ao protocolo de preparo de amostras para MET. Nas imagens obtidas no microscópio Zeiss EM 109, observou-se que os óleos essenciais apresentaram ação fungitóxica notória sobre os conídios de *C. gloeosporioides* e *C. musae*, causando danos severos aos mesmos, promovendo a desorganização e degradação celular que inviabilizaram a germinação.

0.5% observed in previous experiments (data not shown).

Except for the essential oil of *Cinnamomum sp.*, extracted by hydrodistillation in a Clevenger type apparatus, the essential oils were supplied by Chamel Industry and Natural Products Commerce, Paraná State, Brazil in 2008. In test tubes, 0.5ml of suspensions of conidia ( $1 \times 10^{10}$  conidia/ml) prepared in sterile distilled water with 1.0% Tween 20®, were mixed with 0.5ml of the 1.0% essential oils solutions to obtain a final concentration of 0.5%. For the control, 0.5ml of spore suspensions were mixed with 0.5ml of sterile distilled water. The tubes remained under agitation in an Orbital Shaker at 100rpm and an average temperature of 25°C. After 24h, the contents

were transferred to Eppendorf tubes and centrifuged for 3min at 6000rpm. The supernatant was discarded and the masses of conidia fixed in modified Karnovsky fixative, composed of 2.5%glutaraldehyde and 2.5% formaldehyde in 0.05M sodium cacodylate buffer, pH 7.2, plus 0.1M  $\text{CaCl}_2$ , and kept in the refrigerator for 24h (primary fixation). To form pellets, the masses of conidia fixed in Karnovsky were centrifuged for 3min at 6000rpm and the supernatant discarded.

After discarding the supernatant fixative, a 1.0% agarose gel and the pellets were mixed by means of a toothpick and heated at  $\sim 45^\circ\text{C}$ , solidifying instantly. The blocks of agarose, after reduction in size, were washed three times for 10min in sodium 0.05 M cacodylate

buffer for post-fixation in 2%  $\text{OsO}_4$ , in a hood, for 4h. The blocks were then washed three times in distilled water and submitted to contrast *en bloc* in an aqueous solution of 0.5% uranyl acetate overnight in the refrigerator. Afterwards, dehydration of the blocks was achieved in an increasing acetone series of 25, 50, 75, 90 and 100% for 10min each, except the last concentration with  $3 \times 10$ min.

The blocks were subsequently embedded, the acetone being replaced by resin in an increasing gradient, remaining 8h in Spurr resin (30%) and acetone (70%), 8h in Spurr resin (70%) and acetone (30%) and  $2 \times 24$ h in pure Spurr resin, at room temperature. Samples were transferred to silicone molds containing polymerized Spurr

resin in half of their volume, covered with pure Spurr resin to fill the molds, and kept at 70°C for 48h for polymerization. The blocks were trimmed with razor blades to a trapezoidal shape with a cutting surface of appropriate size.

Initially, semithin 0.5 $\mu\text{m}$  sections were made on Reichert-Jung (Ultracut E) ultramicrotome with a glass knife, for the localization of fungal structures of interest in a light microscope and for orientation of the ultrathin sections. The semithin sections were collected with a gold ring and placed on glass slides, dried in a metal plate at  $\sim 60^\circ\text{C}$ , covered with toluidine blue stain (1g toluidine blue, 1g sodium borate and 100ml of water, filtered through a Millipore 0.2 $\mu\text{m}$ ), heated on a

metal plate until formation of a golden border, washed with distilled water, dried on a hot plate and visualized by light microscopy.

Ultrathin sections (>100nm) were made with a diamond knife, collected on copper grids (300 mesh) previously coated with formvar film, post-contrasted with aqueous solutions of 2% uranyl acetate and 3% lead citrate for 3min on each, and washed with distilled water. The observation was carried out with a transmission electron microscope Zeiss EM 109 at 80kV. The images were digitally recorded and edited in the Photopaint Software of the Corel Draw 13 package.

## Results and Discussion

Figures 1a and 2a illustrate the integrity of the cell wall, the plasmatic membrane and cytoplasmic contents of conidia of *Colletotrichum gloeosporioides* and *C. musae*, respectively, in control, untreated preparations.

In the conidia treated with the essential oils of *C. martinii* (Figures 1b and 2b), *E. caryophyllata* (1c and 2c), *T. vulgaris* (1d and 2d), *C. citratus* (1e and 2e) and *Cinnamomum* sp. (Figures 1f and 2f) changes were observed in the cell wall and in the plasmatic membrane, as well as vacuolization of the cytoplasm.

The number of studies demonstrating the mode of action of essential oils on plant pathogenic fungi is small; however, the results mentioned below, demonstrating cellular structural changes observed in other pathosystems and essential oils, corroborate those obtained in the present study.

Zambonelli *et al.* (2004) verified that the essential oil of *T. vulgaris* (thyme), containing




Figure 1. Transmission electron micrographs of untreated conidia of *Colletotrichum gloeosporioides* (control, a), and treated with essential oils of *Cymbopogon martinii* (b); *Eugenia caryophyllata* (c), *Thymus vulgaris* (d), *Cymbopogon citratus* (e) and *Cinnamomum* sp. (f) at 0.5%. CW: cell wall, PM: plasmatic membrane, LB: lipid droplets, V: vacuolization, EM: electron-dense material, and EXT: leakage.

thymol as its main component, caused an increase in the vacuolization of the cytoplasm and an accumulation of lipid droplets, ripples in the plasmalemma and changes in the mitochondria and endoplasmic reticulum of *Colletotrichum lindemuthianum* and *Pythium ultimum*. Rasooli *et al.* (2006) observed severe hyphae collapsing, plasmatic membrane rupture and destruction of mitochondria in *Aspergillus niger* treated with essential oils of *Thymus eriocalyx* and *T. x-parlock*.

The oils of *T. eriocalyx* and *T. x-parlock* were also found to produce irreversible damage to the walls, membranes and cel-

lular organelles by exposing spores of the pathogen. Arroyo *et al.* (2007) found that the volatile compound (E)-hex-2-enal caused changes in the structures of the cell wall and the plasmatic membrane, with consequent disorganization and destruction of organelles and, eventually, cell death of *Colletotrichum acutatum*, one of the agents that cause anthracnose in strawberries.

The accumulation of electron-dense material observed in conidia of *C. gloeosporioides* (Figure 1c) and *C. musae* (Figure 2c) treated with the essential oil of *E. caryophyllata*, characterized by the formation of an electron dark image cor-

roborates the findings of Bakkali (2008), who mentions that, as lipophilic substances, the essential oils penetrate the cell wall and the plasmatic membrane, disrupting the structure of different layers of polysaccharides, fat acids and phospholipids, making them permeable.

The essential oils of *C. citratus* (Figures 1e and 2e) and *Cinnamomum* sp. (Figures 1f and 2f) promoted the leakage of cytoplasmic contents of some conidia. Piper *et al.* (2001) pointed out that substances found in essential oils affect the integrity of cell membranes making them permeable, causing leakage of cellular content.

It was observed that in treatments with the same essential oils occurred variations in ultrastructure of the pathogen's conidia, such as different intensities of vacuolization and leakage of cytoplasmic contents or not. For Knobloch *et al.* (1988) such variations may occur depending on the solubility of essential oils and interaction with

the cytoplasmic membrane, thereby determining the antifungal activity. The thickness of the spores' walls may also interfere with the activity of antifungal compounds, as reported by Svircev *et al.* (2007) who did not find any effect of thymol vapors on the cytoplasm of the thick-walled spores of *Monilinia fructicola* in the postharvest treatment of plum.

From the efficacy of essential oils, demonstrated in the total inhibition of the germination of *C. gloeosporioides* and *C. musae*, it can be inferred that such fungitoxic action occurs on other species of *Colletotrichum*, avoiding the dissemination.

The exploration of antifungal activity of essential oils is presented as an alternative strategy for the control of plant diseases in pre-and post-harvest, representing a lesser risk to human health and environment in pre-and post-harvest and could replace pesticides.

## Conclusion

The essential oils of *Cinnamomum* sp., *Cymbopogon citratus*, *Eugenia caryophyllata*, *Cymbopogon martinii* and *Thymus vulgaris* presented direct fungitoxic action on *C. gloeosporioides* and *C. musae*, causing severe damage to cellular ultrastructure of the conidia.

## ACKNOWLEDGEMENTS

The authors acknowledge the doctoral scholarship and financial support granted by the National Council of Scientific and Technological Development (CNPq), and the financial support of the Research Foundation of the State of Minas Gerais (FAPEMIG) to the Laboratory of Electron Microscopy and Ultrastructural Analysis at the Federal University of Lavras, MG, Brazil.

## REFERENCES

Anaruma ND, Schmidt FL, Duarte MCT, Figueira GM, Delarmelina C, Benato EA, Sartoratto A (2010) Control of *Colletotrichum gloeosporioides* (penz.) Sacc. in yellow passion fruit using *Cymbopogon citratus* essential oil. *Braz. J. Microbiol.* 41: 66-73.

Arroyo FT, Moreno J, Daza P, Boianova L, Romero F (2007) Antifungal activity of strawberry fruit volatile compounds against *Colletotrichum acutatum*. *J. Agric. Food Chem.* 55: 5701-5707.

Bakkali F, Averbeck S, Averbeck D, Idaomar M (2008) Biological




Figure 2. Transmission electron micrographs of untreated conidia of *Colletotrichum musae* (control, a), and treated with essential oils of *Cymbopogon martinii* (b); *Eugenia caryophyllata* (c), *Thymus vulgaris* (d), *Cymbopogon citratus* (e) and *Cinnamomum* sp. (f) at 0.5%. CW: cell wall, PM: plasmatic membrane, LB: lipid droplets, V: vacuolization, EM: electron-dense material, and EXT: leakage.

effects of essential oils: a review. *Food Chem. Toxicol.* 46: 446-475.

Barrera-Necha LL, Bautista-Baños S, Flores-Moctezuma HE, Rojas-Estudillo A (2008) Efficacy of essential oils on the conidial germination, growth of *Colletotrichum gloeosporioides* (Penz.) Penz. and Sacc and control of postharvest diseases in papaya (*Carica papaya* L.). *Plant Pathol.* 7: 174-178.

Duamkhanmanee R (2008) Natural essential oils from lemon grass (*Cymbopogon citratus*) to control postharvest anthracnose of mango fruit. *Int. J. Biotechnol.* 10: 104-108.

Knobloch K, Pauli A, Iberl B, Weis N, Weigand H (1988) Mode of action of essential oil components on whole cells and fungi in plate tests. Walter de

Gruyter. Berlin, Germany. pp. 287-299.

Lee SO, Choi GJ, Jang KS, Lim HK, Cho KY, Kim JC (2007) Antifungal activity of five plant essential oils as fumigant against postharvest and soilborne plant pathogenic fungi. *Plant Pathol.* 23: 97-102.

Peres NAR, Kuramae EE, Dias MSC, Souza NL de (2002) Identification and characterization of *Colletotrichum* spp. affecting fruit after harvest in Brazil. *J. Phytopathol.* 150: 128-134.

Piper P, Calderón CO, Hatzixanthos K, Mollapour M (2001) Weak acid adaptation: the stress response that confers yeasts with resistance to organic acid food preservatives. *Microbiology* 147: 2635-2642.

Ranasinghe L, Jayawardena B, Abeywickrama K (2002) Fungicidal activity of essential oils of

*Cinnamomum zeylanicum* (L.) and *Syzygium aromaticum* (L.) Merr et L.M. Perry against crown rot and anthracnose pathogens isolated from banana. *Lett. Appl. Microbiol.* 35: 208-211.

Rasooli I, Rezaei MB, Allameh A (2006) Growth inhibition and morphological alterations of *Aspergillus niger* by essential oils from *Thymus eriocalyx* and *Thymus x-porlock*. *Food Control* 17: 359-364.

Rozwalka LC, Lima ML-RZC, May-De Mio LL, Nakashima T (2008) Extratos, decoctos e óleos essenciais de plantas medicinais e aromáticas na inibição de *Glomerella cingulata* e *Colletotrichum gloeosporioides* de frutos de goiaba. *Ciência Rural* 38: 31-36.

Shahi SK, Patra M, Shukla AC, Dikshit A (2003) Use of essential oil as botanical-pesticide against post harvest spoilage in *Malus pumilo* fruit. *BioControl* 48: 223-232.

Siani AC, Sampaio ALF, Sousa MC de, Henriques M das GMO, Ramos MFde (2000) Óleos essenciais: potencial anti-inflamatório. *Rev. Biotechnol. Ciênc. Desenv.* 3: 38-43.

Singh G, Singh OP, Rao GP, Sharma SR (2002) The vapour action of essential oils and monoterpenoids against pathogenic fungi. *Sugar Technol. J.* 4: 69-71.

Sukatta U, Haruthaithanasan V, Chantarapanont W, Dilokkunanant U, Suppakul P (2008) Antifungal activity of clove and cinnamon oil and their synergistic against postharvest decay fungi of grape *in vitro*. *Nat. Sci.* 42: 169-174.

Svircev AM, Smith RJ, Zhou T, Hernández M, Liu W, Chud CL (2007) Effects of timol fumigation on survival and ultrastructure of *Monilinia fructicola*. *Postharv. Biol. Technol.* 45: 228-233.

Tzortzakakis NG (2009) Impact of cinnamon oil-enrichment on microbial spoilage of fresh produce. *Innov. Food Sci. Emerg. Technol.* 10: 97-102.

Tzortzakakis NG, Economakis CD (2007) Antifungal activity of lemongrass (*Cymbopogon citratus* L.) essential oil against key postharvest pathogens. *Innov. Food Sci. Emerg. Technol.* 8: 253-258.

Zambonelli A, D'aulerio AZ, Severi A, Benvenuti S, Maggi L, Bianchi A (2004) Chemical composition and fungicidal activity of commercial essential oils of *Thymus vulgaris* L. *J. Essent. Oil Res.* 16: 69-74.